Mark scheme

Logarithmic and exponential functions

Additional Mathematics 0606 · Questions from Paper 1 and Paper 2

Total marks

71

MARKS FOR EACH QUESTION

1	2	3	4	5	6	7	8	9	10	Total
6	4	5	7	7	7	8	8	9	10	**71**

Question 1 [6]

1(i)	The number of bacteria at the start of the experiment	**B1**	
1(ii)	$20\,000 = 800e^{kt}$ so $\frac{20\,000}{800} = e^{2k}$ or $\ln 20\,000 = \ln 800 + \ln(e^{2k})$	**M1**	use of given equation and attempt to solve for e^{2k} or use logs correctly
	$2k = \ln 25$	**M1**	correct method to obtain $2k$
	1.61	**A1**	
1(iii)	$P = 800e^{3\ln 5}$	**M1**	Substitution of $t = 3$ in formula using *their k*
	$= 100\,000$	**A1**	answer in range 99800 to 100200

Question 2 [4]

2	$4\times 8^{2x} + 8^{x} - 3[=0]$ or $4\times(2^{3x})^{2} + 2^{3x} - 3[=0]$ oe	**B1**	
	$\left(4(8^{x}) - 3\right)\left(8^{x} + 1\right)$ or $\left(4(2^{3x}) - 3\right)\left(2^{3x} + 1\right)$ oe	**M1**	**FT** *their* 3-term quadratic in 8^{x} or 2^{3x}
	$8^{x} = \frac{3}{4}$ or $2^{3x} = \frac{3}{4}$ oe, nfww	**A1**	
	$x = \log_{8}(0.75)$ or $\frac{1}{3}\log_{2}(0.75)$ or $-0.138[34\ldots]$ oe	**A1**	and no other solutions

STEM | Exam Builder

Question 3 [5]

3	Correct change of base e.g. $\log_2(x+1)-\dfrac{4\log_2 2}{\log_2(x+1)}=3$ or $\dfrac{\log_{(x+1)}(x+1)}{\log_{(x+1)}2}-4\log_{(x+1)}2=3$	**B1**	
	$\left(\log_2(x+1)\right)^2-3\log_2(x+1)-4 \quad [=0]$ oe or $4\left(\log_{(x+1)}2\right)^2+3\log_{(x+1)}2-1 \quad [=0]$ oe	**B1**	
	Factorises or solves e.g. $\left(\log_2(x+1)-4\right)\left(\log_2(x+1)+1\right)=0$ or $\left(4\log_{(x+1)}2-1\right)\left(\log_{(x+1)}2+1\right)=0$	**M1**	**FT** *their* 3-term quadratic in a suitable logarithm providing correct change of base seen
	$\log_2(x+1)=their 4$ and $\log_2(x+1)=their(-1)$ or $\log_{(x+1)}2=their\dfrac{1}{4}$ and $\log_{(x+1)}2=their(-1)$ **and** correctly solves as far as x = … at least once	**M1**	**dep** on previous **M1** **FT** $\log_2(x+1)=a$ or $\log_{(x+1)}2=b$ where a and b are constants
	$x=15,\ x=-\dfrac{1}{2}$ nfww mark final answer	**A1**	

Question 4 [7]

4 **(i)**	$2000=1000e^{a+b} \rightarrow a+b=\ln 2$	**B1**	
(ii)	$3297=1000e^{2a-b} \rightarrow 2a+b$ $=\ln 3.297$ oe	**M1** **A1**	substitution of 2, 3297 and rearrange
(iii)	Solve for one value $a=0.5$ and $b=0.193$ or 0.19	**M1** **A1**	
(iv)	$n=10 \quad P=1000e^{5.193}$ $=\$180\,000.$	**M1** **A1**	

STEM | Exam Builder

Question 5 **[7]**

5(a)	Either $2\log_4 y = \log_2 y$ Or $\log_2 x = 2\log_4 x$	**B1**	
	Either $\log_2 x + \log_2 y = 8$ leading to $\log_2 xy = 8$ Or $2\log_4 x + 2\log_4 y = 8$ leading to $\log_4 xy = 4$	**M1**	For use of log law
	$xy = 256$	**A1**	
5(b)	$2y^2 - 3y + 1 = 0$	**B1**	
	$y = \frac{1}{2},\ 1$	**M1**	For attempt to solve for y
	$x = -1$	**A1**	
	$x = 0$	**A1**	

STEM | Exam Builder

Question 6 **[7]**

6(a)	$(2y-1)\log 5 = \log 6 + y\log 3$ oe or $2y\log 5 = \log 30 + y\log 3$ oe OR [rearranges $\frac{5^{2y}}{5} = 6\times 3^{y}$ and collects powers to a single power in y] $\left(\frac{5^2}{3}\right)^{y} = 30$ oe	**M1**	
	Collects terms and factorises: $y(2\log 5 - \log 3) = \log 6 + \log 5$ oe or $y(2\log 5 - \log 3) = \log 30$ oe OR takes logs $y = \log_{\frac{25}{3}} 30$ oe or $y\log\left(\frac{5^2}{3}\right) = \log 30$ oe	**M1**	**FT** if of equivalent difficulty
	1.604	**A1**	cao

6(b)	$e^{4x} - 4e^{2x} + 3 = 0$ or $(e^{2x})^2 - 4e^{2x} + 3 = 0$ oe	**M1**	condone one error
	Factorises: $(e^{2x}-1)(e^{2x}-3)$ oe or solves $e^{4x} - 4e^{2x} + 3 = 0$ oe	**M1**	**FT** $a(e^{2x})^2 + be^{2x} + c[=0]$
	$e^{2x} = 1, \quad e^{2x} = 3$	**A1**	
	$x = 0, \ x = \frac{1}{2}\ln 3$ or exact equivalent	**A1**	

STEM | Exam Builder

Question 7 **[8]**

7(a)	$2x-3=6^{\frac{1}{2}}$ oe, soi	**M1**	
	$x=\dfrac{6^{\frac{1}{2}}+3}{2}$ or $x=\dfrac{\sqrt{6}+3}{2}$	**A1**	
7(b)	$\ln\dfrac{2u}{u-4}=\ln \mathrm{e}$ soi or $\ln\dfrac{2u}{u-4}=1$ soi or $\ln 2u=\ln \mathrm{e}(u-4)$ soi	**M1**	Condone one sign or bracketing error
	$\dfrac{2u}{u-4}=\mathrm{e}$ or $2u=\mathrm{e}(u-4)$ oe	**M1**	**FT** *their* logarithmic equation
	$u=\dfrac{4\mathrm{e}}{\mathrm{e}-2}$ or $u=\dfrac{-4\mathrm{e}}{2-\mathrm{e}}$ or equivalent exact form	**A1**	
7(c)	$\dfrac{3^v}{\left(3^3\right)^{2v-5}}=3^2$ oe soi or $\dfrac{9^{\frac{v}{2}}}{\left(9^{\frac{3}{2}}\right)^{2v-5}}=9$ oe soi or $\log 3^v-\log 27^{2v-5}=\log 9$ oe soi	**B1**	
	$15-5v=2$ oe or $v\log 3-(2v-5)\log 27=\log 9$	**M1**	**FT** *their* exponential equation in the same base or *their* logarithmic equation with any consistent base, providing *their* exponential or logarithmic equation has at most one sign or arithmetic error
	$v=\dfrac{13}{5}$ oe	**A1**	

STEM | Exam Builder

Question 8 **[8]**

8(a)	$\frac{3^{10x}}{3^{3x-6}}[=243]$ oe or $\log 9^{5x} - \log 27^{x-2} = \log 243$ oe	**B1**	
	$3^{7x+6} = 3^5$ soi oe or $5x(\log 9) - (x-2)\log 27 = \log 243$	**M1**	
	$x = -\frac{1}{7}$	**A1**	

8(b)	$\frac{1}{2}\log_a b - \frac{1}{2} = \frac{1}{\log_a b}$ or $\frac{\frac{1}{2}}{\log_b a} - \frac{1}{2} = \log_b a$	**B2**	**B1** for bringing down the power of $\frac{1}{2}$ e.g. $\frac{1}{2}\log_a b$ or for a change of base e.g. $\frac{1}{\log_a b}$
	Clears the fraction and rearranges $\frac{1}{2}(\log_a b)^2 - \frac{1}{2}\log_a b = 1$ oe $(\log_a b)^2 - \log_a b - 2 = 0$ oe or let $x = \log_a b$ $x^2 - x - 2 = 0$ oe or $\frac{1}{2} - \frac{1}{2}\log_b a = (\log_b a)^2$ $0 = 2(\log_b a)^2 + \log_b a - 1$ oe or let $y = \log_b a$ $2y^2 + y - 1 = 0$	**M1**	
	$(\log_a b - 2)(\log_a b + 1)$ oe or $(2\log_b a - 1)(\log_b a + 1)$	**M1**	
	[$\log_a b = 2$, $\log_a b = -1$ or $\log_b a = \frac{1}{2}$, $\log_b a = -1$ leading to] $b = a^2$, $b =$ oe	**A1**	

STEM | Exam Builder

Question 9 [9]

9(a)	$\lg 100 = 2$	**B1**	
	$3\lg x = \lg x^3$	**B1**	
	$\lg \dfrac{100x^3}{y}$	**B1**	
9(b)(i)	$6x^2 + 7x - 3 = 0$ $(2x+3)(3x-1) = 0$	**M1**	For obtaining in suitable quadratic form and attempt to solve
	$x = -\dfrac{3}{2}$ $x = \dfrac{1}{3}$	**A1**	For both
9(b)(ii)	$x = \log_a 3$ $\dfrac{1}{3} = \log_a 3$	**M1**	For realising connection with part **(i)** and attempt to solve $\dfrac{1}{3} = \log_a 3$ or $-\dfrac{3}{2} = \log_a 3$
	$a = 27$	**A1**	
	$-\dfrac{3}{2} = \log_a 3$	**M1**	For realising connection with part **(i)** and attempt to solve $-\dfrac{3}{2} = \log_a 3$ or $\dfrac{1}{3} = \log_a 3$
	$a = \left(\dfrac{1}{3}\right)^{\frac{2}{3}}$ or 0.481or $\left(\dfrac{1}{9}\right)^{\frac{1}{3}}$ oe	**A1**	

Question 10 [10]

10 **(a)(i)** Express in powers of 5. $\left(\dfrac{5^{2x+3}}{5^{4x}} = \dfrac{5^{2(2-x)}}{5^{3x}}\right)$		M1
Use rules of indices ($2x + 3 - 4x = 2(2 - x) - 3x$)		M1
$\dfrac{1}{3}$		A1
(ii) LHS = $\lg y(y - 15)$		B1
$2 = \lg 100$		B1
Solve 3 term quadratic		M1
20 only		A1
(b) $\log_{12} 16 - \log_{12} 9 + \log_{12} 81$		B1
Correctly combine 3 logarithms		M1
2		A1
		[10]

STEM | Exam Builder